

IC/94/65
Napoli: DSF-T-5/94, INFN-NA-IV-5/94
March 1994

Z FLUX-LINE LATTICES AND SELF-DUAL EQUATIONS IN THE STANDARD MODEL

G. Bimonte

and

G. Lozano ¹

*International Centre for Theoretical Physics, P.O.BOX 586
I-34100 Trieste, ITALY*

Abstract

We derive gauge covariant self-dual equations for the $SU(2) \times U(1)_Y$ theory of electro-weak interactions and show that they admit solutions describing a periodic lattice of Z-strings.

¹ E-mail addresses: Bimonte@ictp.trieste.it , Lozano@ictp.trieste.it

Solutions to the classical equations of motion of the Standard Model have recently attracted a considerable attention after the observation made by Vachaspati [1] that the embedding [2] of the Nielsen-Olesen [3] string into the Standard Model can be classically stable in a certain range of parameters. Further studies [4] have revealed that the region of classical stability does not overlap with the one compatible with experimental data. It has been suggested [5] that the cause of this instability might be traced back to the phenomenon of W-condensation [6] but the search of stable W-dressed strings has given negative results [7].

In their original investigations on electroweak magnetism, Ambjørn and Olesen [6] have solved the electroweak equations showing the existence of classically stable configurations representing a condensate of W bosons. Classical stability is ensured since these configurations satisfy a set of first order differential equations (self-dual equations) which arise from the requirement that they saturate a Bogomol'nyi like bound [8] for the energy. The solutions found in Ref. [6] carry no Z flux. The reason for this is that the self-dual equations were derived in the unitary gauge which excludes from the beginning the existence of Z -strings.

In this work we present a *gauge covariant* expression of the self-dual equations and show that there exist boundary conditions which lead to configurations carrying Z flux.

We will consider static configurations which are axially symmetric along the z -axis and such that physical observables are periodic in the $x - y$ plane. The energy functional for a cell $\mathcal{C}$ of periodicity with area A is then given by

$$E = \int_{\mathcal{C}} d^2x \left[\frac{1}{4} W_{ij}^a W_{ij}^a + \frac{1}{4} B_{ij} B_{ij} + (D_i \Phi)^\dagger D_i \Phi + \lambda (\Phi^\dagger \Phi - \phi_0^2)^2 \right] , \quad (1)$$

where W_{ij}^a and B_{ij} are the field strengths associated with the gauge group $SU(2) \times U(1)_Y$

$$W_{ij}^a = \partial_i W_j^a - \partial_j W_i^a + g \epsilon_{abc} W_i^b W_j^c ,$$

$$B_{ij} = \partial_i B_j - \partial_j B_i , \quad (2)$$

while the covariant derivative for the Higgs-doublet Φ is defined as

$$D_i \Phi = \left(\partial_i - \frac{i}{2} g \tau^a W_i^a - \frac{i}{2} g' B_i \right) \Phi. \quad (3)$$

In Eq.(3), τ^a are the Pauli matrices and g and g' are related to the Weinberg angle θ_W and to the electric charge e by the relations

$$e = g \sin \theta_W \quad g' = g \tan \theta_W. \quad (4)$$

The electromagnetic field A_i , the neutral Z_i field and the charged W_i field are defined as:

$$\begin{aligned} A_i &= \sin \theta_W W_i^3 + \cos \theta_W B_i \quad , \\ Z_i &= \cos \theta_W W_i^3 - \sin \theta_W B_i \quad , \\ W_i &= \frac{1}{\sqrt{2}}(W_i^1 - iW_i^2) \quad . \end{aligned} \quad (5)$$

and the masses of the physical particles are given by:

$$m_W^2 = \frac{1}{2}g^2\phi_0^2 \quad , \quad m_Z^2 = \frac{g^2\phi_0^2}{2\cos^2\theta_W} \quad , \quad m_H^2 = 4\lambda\phi_0^2 \quad . \quad (6)$$

In order to find a bound for the energy, we will start by rewriting the kinetic energy of the Higgs field in an appropriate way by extending to the electroweak theory the identity used by Bogomol'nyi in the simpler $U(1)$ Landau-Ginzburg theory [8]; this identity reads:

$$\begin{aligned} |D_i\Phi|^2 &= \frac{1}{2}|D_i\Phi - i\gamma\epsilon_{ij}D_j\Phi|^2 + \\ &\quad -\frac{1}{4}\gamma g\Phi^\dagger\tau^a\Phi\epsilon_{ij}W_{ij}^a - \frac{1}{4}\gamma g'\Phi^\dagger\Phi\epsilon_{ij}B_{ij} + \gamma\epsilon_{ij}\partial_i J_j \end{aligned} \quad (7)$$

where

$$J_j = \frac{1}{2i}[\Phi^\dagger D_j\Phi - (D_j\Phi)^\dagger\Phi] \quad (8)$$

and $\gamma^2=1$ In what follows we will set $\gamma = -1$ (the opposite case can be treated analogously). After making use of Eq.(7), the energy can be written as:

$$E = E_1 + \frac{g'\phi_0^2}{4\sin^2\theta_W} \int_{\mathcal{C}} d^2x \epsilon_{ij} B_{ij} - \frac{g^2\phi_0^4}{8\sin^2\theta_W} \int_{\mathcal{C}} d^2x + \left(\lambda - \frac{g^2}{8\cos^2\theta_W} \right) \int_{\mathcal{C}} d^2x (\Phi^\dagger \Phi - \phi_0^2)^2 +$$

$$- \int_{\mathcal{C}} d^2x \epsilon_{ij} \partial_i J_j \quad , \quad (9)$$

where E_1 is the positive definite functional

$$E_1 = \int_{\mathcal{C}} d^2x \left\{ \frac{1}{4} \left[W_{ij}^a + \epsilon_{ij} \frac{g}{2} \Phi^\dagger \tau^a \Phi \right]^2 + \frac{1}{4} \left[B_{ij} + \epsilon_{ij} \frac{g'}{2} \left(\Phi^\dagger \Phi - \frac{\phi_0^2}{\sin^2\theta_W} \right) \right]^2 + \right.$$

$$\left. + \frac{1}{2} |D_i \Phi + i\epsilon_{ij} D_j \Phi|^2 \right\} . \quad (10)$$

The last term in Eq.(9) can always be dropped after integration on a cell, since it is the divergence of a gauge invariant quantity (remember that observables are periodic). The second term in (9) is of topological nature and is proportional to the flux Φ_B of the $U(1)_Y$ field B_i , while the third term is a geometric term proportional to the area A of the cell. We then see from (9) that for

$$\lambda \geq \frac{g^2}{8\cos^2\theta_W} \quad (11)$$

or equivalently

$$m_H \geq m_Z \quad , \quad (12)$$

the energy is bounded by

$$E_{\text{bound}} = \frac{g'\phi_0^2}{2\sin^2\theta_W} \Phi_B - \frac{g^2\phi_0^4}{8\sin^2\theta_W} A = \frac{m_W^2}{2e^2} (2g'\Phi_B - m_W^2 A) . \quad (13)$$

This is the Bogomol'nyi bound for the Standard Model, derived in a gauge covariant way. The simpler $U(1)$ case is recovered by setting $g = 0$ in Eq.(13). A significant difference between the abelian and the non-abelian case is the presence in the latter of the term proportional to the area, which spoils the topological nature of the bound. Although (12) covers most of the experimental region ($m_H > 0.62m_Z$), let us point out that a slight modification of (7) as in Ref. [9] allows one to deduce an expression of the bound valid in the case $m_H < m_Z$:

$$E_{\text{bound}} = \frac{1}{2e^2} \left(\frac{m_W m_H}{m_Z} \right)^2 \left[2g'\Phi_B - \left(\frac{m_W m_H}{m_Z} \right)^2 A \right] . \quad (14)$$

Now, let us concentrate on configurations saturating the bound. This occurs only if $m_H = m_Z$ and $E_1 = 0$, which in turn implies:

$$W_{ij}^a = -\epsilon_{ij} \frac{g}{2} \Phi^\dagger \tau^a \Phi \quad , \quad (15)$$

$$B_{ij} = -\epsilon_{ij} \frac{g'}{2} \left(\Phi^\dagger \Phi - \frac{\phi_0^2}{\sin^2 \theta_W} \right) \quad , \quad (16)$$

$$D_i \Phi = -i\epsilon_{ij} D_j \Phi \quad . \quad (17)$$

This is the gauge covariant form of the self-dual (or Bogomol'nyi) equations previously found in the unitary gauge by Ambjørn and Olesen [6].

Of course, for $g = 0$, Eqs.(15-17) become the well studied self-dual equations of the $U(1)$ model [8, 10, 11]. For the purely non-abelian case, which is obtained by taking $g' = 0$, it is in some way surprising that the same equations appear in a quite different context, that of a theory of non-relativistic bosons interacting with a Chern-Simons field [12].

We will now look for solutions of (15-17), corresponding to periodic observables in the plane. In order to simplify the discussion of boundary conditions it is convenient to point out that every solution of Eqs. (15-17) can be converted via a smooth gauge transformation into the *generalized* unitary gauge

$$\Phi = \begin{pmatrix} 0 \\ \phi \end{pmatrix} \quad . \quad (18)$$

where ϕ is *complex*. This follows from the fact that all Higgs doublets satisfying the Bogomol'nyi Eq.(17) can be written locally as:

$$\Phi = (z - z_0)^n \Psi(x, y) \quad , \quad (19)$$

where $z = x + iy$ and $\Psi(x, y)$ is a doublet such that:

$$(\Psi^\dagger \Psi)(z_0) \neq 0.$$

It then follows that Ψ can be converted to the unitary gauge locally. This transformation can always be extended globally to the plane without obstructions. The most general boundary conditions which give periodic observables are then (for simplicity we consider a rectangular standard cell of sides L_X and L_Y):

$$\Phi(x + L_X, y) = e^{i[\alpha_X(x, y)\tau_3 + \beta_X(x, y)]} \Phi(x, y)$$

$$\begin{aligned}
W_i(x + L_X, y) &= e^{2i\alpha_X(x, y)} W_i(x, y) \\
W_i^3(x + L_X, y) &= W_i^3(x, y) + \frac{2}{g} \partial_i \alpha_X(x, y) \\
B_i(x + L_X, y) &= B_i(x, y) + \frac{2}{g'} \partial_i \beta_X(x, y),
\end{aligned} \tag{20}$$

with analogous boundary conditions in the y direction, with functions α_Y and β_Y . It is easily verified that α_Y and β_Y can be gauged away and we can then assume without any loss of generality that all fields are periodic in the y direction. Consistency of the boundary conditions (20) at the corners of the cell implies the following constraints for α_X and β_X :

$$\begin{aligned}
\alpha_X(x, y) &= \frac{\pi k_3}{L_Y} y + \tilde{\alpha}_X(x, y) \\
\beta_X(x, y) &= \frac{\pi k}{L_Y} y + \tilde{\beta}_X(x, y) ,
\end{aligned} \tag{21}$$

where k and k_3 are integers such that $k - k_3$ is even. The functions $\tilde{\alpha}_X$ and $\tilde{\beta}_Y$ are periodic in y and can be gauged away. Summarizing, the most general boundary conditions compatible with Eqs. (15-17) read:

$$\begin{aligned}
\phi(x + L_X, y) &= e^{i\pi(k - k_3) \frac{y}{L_Y}} \phi(x, y) , \\
W_i(x + L_X, y) &= e^{2i\pi k_3 \frac{y}{L_Y}} W_i(x, y) , \\
W_i^3(x + L_X, y) &= W_i^3(x, y) + \frac{2\pi k_3}{g L_Y} \delta_{i,2} , \\
B_i(x + L_X, y) &= B_i(x, y) + \frac{2\pi k}{g' L_Y} \delta_{i,2} .
\end{aligned} \tag{22}$$

Boundary conditions (22) reduce to those considered in [6] only if $k = k_3$. In this case, ϕ is real and periodic. Solutions with different values of k and k_3 are gauge inequivalent.

The integers k and k_3 are related to the fluxes of f_{12} and Z_{12} :

$$\Phi_A \equiv \int_C d^2 x f_{12} = \frac{2\pi k}{e} - \frac{2\pi \sin^2 \theta_W}{e} (k - k_3) , \tag{23}$$

$$\Phi_Z \equiv \int_C d^2 x Z_{12} = -\frac{\pi}{e} \sin 2\theta_W (k - k_3) . \tag{24}$$

Using Eq.(18) in Eq. (17), we now obtain the following equation for the field W_i :

$$(W_i + i\epsilon_{ij} W_j) \phi = 0 , \tag{25}$$

which implies that, outside the zeroes of ϕ ,

$$W_1 = -iW_2 \equiv W \quad . \quad (26)$$

Using this relation and Eq. (18) into eqs (15-17), we get:

$$\tilde{D}_i W + i\epsilon_{ij} \tilde{D}_j W = 0 \quad , \quad (27)$$

$$D_i \hat{\phi} + i\epsilon_{ij} D_j \hat{\phi} = 0 \quad , \quad (28)$$

$$f_{12} = \frac{m_W^2}{e} + 2e|W|^2 \quad , \quad (29)$$

$$Z_{12} = \frac{m_W^2 \tan \theta_W}{e} (\hat{\phi}^* \hat{\phi} - 1) + \frac{2e}{\tan \theta_W} |W|^2 \quad , \quad (30)$$

where $\hat{\phi} = \phi/\phi_0$ and

$$\tilde{D}_i = \partial_i - ieA_i - i\frac{e}{\tan \theta_W} Z_i \quad ,$$

$$D_i = \partial_i + i\frac{e}{\sin 2\theta_W} Z_i \quad .$$

Eq. (27-30) have solutions only when k_3 and $k - k_3$ are non-negative (the opposite statement holds when $\gamma = 1$). Notice that the energy of configurations satisfying the Bogomol'nyi equations depends only on k ,

$$E = \frac{\phi_0^2}{\sin^2 \theta_W} \pi k - \frac{g^2 \phi_0^4}{8 \sin^2 \theta_W} A \quad . \quad (31)$$

We then see that different values of k_3 correspond to degenerate but not gauge equivalent solutions.

These equations are consistent only if the area A of the standard cell belongs to a range determined by the fluxes and the parameters of the theory. By integrating Eq. (29), we get:

$$e\Phi_A = m_W^2 A + 2e^2 \int_C d^2 x |W|^2 \geq m_W^2 A \quad , \quad (32)$$

or

$$m_W^2 A \leq 2\pi k - 2\pi \sin^2 \theta_W (k - k_3) \quad . \quad (33)$$

The lower bound on the area can be obtained directly from Eq.(16):

$$g' \Phi_B = -g'^2 \int_C d^2 x \Phi^\dagger \Phi + \frac{g'^2 \phi_0^2}{2 \sin^2 \theta_W} A \leq \frac{g'^2 \phi_0^2}{2 \sin^2 \theta_W} A \quad , \quad (34)$$

or

$$m_W^2 A \geq 2\pi k \cos^2 \theta_W \quad . \quad (35)$$

These bounds coincide with those derived in [6] when $k = k_3$. Notice nevertheless that while the upper bound still depends only on the flux of f_{12} , the lower bound now also depends on the flux of Z_{12} . It is worth to analyze the case when the area A is equal to the upper bound (33). In this case, we obtain a solution by setting

$$W = 0 \quad ,$$

$$f_{12} = \frac{m_W^2}{e} \equiv H_{c_1} \quad . \quad (36)$$

The equations for $\hat{\phi}$ and Z_i become:

$$D_i \hat{\phi} + i\epsilon_{ij} D_j \hat{\phi} = 0 \quad , \quad (37)$$

$$Z_{12} = \frac{m_W^2 \tan \theta_W}{e} (\hat{\phi}^* \hat{\phi} - 1) \quad . \quad (38)$$

Eqs. (37) and (38) are the self-dual equations of a $U(1)$ theory defined on a torus. Although no explicit form of the solutions is available, their existence as well as the properties of the associated moduli space have been studied in detail [13]. Notice nevertheless the presence of a constant magnetic field (36) which is the remnant of the non-abelian structure of the equations. This shows itself also in the expression of the energy of these configurations which differs from the abelian formula by a term proportional to the area, which is precisely the energy of the electromagnetic field,

$$E = \frac{e\phi_0^2}{\sin 2\theta_W} |\Phi_Z| + \frac{1}{2} H_c^2 A \quad . \quad (39)$$

We have then derived a gauge covariant expression for the self-dual equations of the Standard Model and showed, in a particular case, the existence of classically stable configurations carrying Z flux. As in Ref. [1], these configurations correspond to an embedding of $U(1)$ into $SU(2) \times U_Y(1)$, but in this case classical stability holds for values of the parameters not ruled out by experiments ($m_H = m_Z$). Finally, our configurations present a superimposed constant magnetic field, a phenomenon that can be linked to the non-abelian nature of the equations. Notice also in this respect that our solutions correspond to a minimization of the energy on a *compact* region.

The structure represented by Eqs. (36-39) is simple but is far from being general. It would be interesting to explicitly show the existence of solutions for $f_{12} > H_{c1}$, in which case a W -condensate will be present. In this regard, we believe that the methods used in Ref. [14] can be applied after suitable modifications.

We would like to thank Prof. Abdus Salam, the International Atomic Energy Agency and UNESCO for hospitality at the International Centre for Theoretical Physics.

References

- [1] T.Vachaspati, Phys. Rev. Lett. **68**, 1997 (1992).
- [2] Y.Nambu, Nucl. Phys. **B130**, 505 (1977);
M.B.Einhorn and A.Savit, Phys. Lett. **77 B**, 295 (1978);
N.S.Manton, Phys. Rev. **D28**, 2018 (1983).
- [3] H.B.Nielsen and P.Olesen, Nucl. Phys. **B61**, 45 (1973).
- [4] T.Vachaspati, Nucl. Phys. **B397**, 648 (1993);
M.James, L.Perivolaropoulos and T.Vachaspati, Nucl. Phys. **B395**, 534 (1993).
- [5] W.B.Perkins, Phys. Rev. **D47**, 5224 (1993);
P.Olesen, Niels Bohr Institute preprint, NBI-HE-93-58, hep-ph/9310275.
- [6] J.Ambjørn and P.Olesen, Nucl. Phys. **B315**, 606 (1989); Nucl. Phys. **B330**, 193 (1990); Int. J. Mod. Phys.**A5**, 4525 (1993).
- [7] A.Achúcarro, R.Gregory, J.A.Harvey and K.Kuijken, E.Fermi Institute preprint, EFI-93-68, hep-th/9312034.
- [8] E.B.Bogomol'nyi, Sov. Jour. Nucl. Phys. **24**,449 (1976);
H. de Vega and F.Schaposnik, Phys. Rev. **D14**, 1160 (1976).
- [9] L.Cugliandolo, G.Lozano and F.Schaposnik, Phys. Rev. **D40**, 3430 (1989).
- [10] A.Jaffe and C.Taubes, *Vortices and Monopoles*, Progress in Physics Vol. 2 (Birkhäuser, Boston, Basel, Stuttgart 1980).
- [11] E.Weinberg, Phys.Rev. **D19**, 3008 (1979)
- [12] G.Dunne, R.Jackiw, S.-Y.Pi and C.A.Trugenberger, Phys. Rev. **D43**, 1332 (1991).
- [13] P.A.Shah and N.S.Manton, DAMTP 93-30.

[14] J.Spruck and Y.Yang, Comm. Math. Phys. **144**, 16 (1992).